



International
Animal Health Products
THE AUSTRALASIAN COMPANY

LIVAMOL[®] with BioWorma[®]

NUTRITIONAL SUPPLEMENT WITH THE NATURAL BIOLOGICAL CONTROL, *BIOWORMA*[®], TO REDUCE INFECTIVE NEMATODE LARVAE WITHIN THE MANURE OF GRAZING ANIMALS

PRESENTATION: A highly palatable, green free flowing fine meal.

ACTIVE CONSTITUENTS: Each kg contains:

VITAMIN A	60,000 I.U.	COPPER	15.7 mg
VITAMIN D ₃	12,000 I.U.	COBALT	80 µg
VITAMIN B ₁	1 mg	IODINE	87 µg
VITAMIN B ₂	2 mg	MANGANESE	104 mg
VITAMIN B ₆	1 mg	ZINC	25 mg
VITAMIN B ₁₂	32 µg		
<i>DUDDINGTONIA FLAGRANS</i> IAH 1297		30 x 10 ⁶ Chlamydospores	

(These levels of active constituents are minimum levels)

NUTRITIONAL INFORMATION:

PROTEIN	Min. 20 %
FAT	Min. 5 %
FIBRE	Min. 9 %
CALCIUM (Ca)	Min. 4 % - Max. 6 %
PHOSPHORUS (P)	Min. 1.5 % - Max. 2.5 %
SALT	NIL
MAXIMUM FLUORINE (F)	0.025 %
Average Digestible Energy	11 MJ/kg

PROPERTIES: *LIVAMOL*[®] with *BioWorma*[®] contains the spores of *Duddingtonia flagrans*, a natural fungus found in soil and on pasture. It is a non-chemical biological control for the free-living stages of parasitic gastrointestinal nematodes of grazing animals, which acts by substantially reducing the numbers of infective worm larvae (including chemical/multi-resistant larvae) emerging from manure onto pasture. When fed to animals, the thick-walled spores remain inert (having no effect within the host animal) and resist digestion, passing through into the manure. There they germinate and form trapping organs that capture, paralyse and consume emerging worm larvae (including chemical/multi-resistant larvae), interrupting the crucial re-infestation stage of the parasites' life cycle, thus reducing the amount of re-infection on the pasture. The spores are safe, non-toxic and residue-free. There are no negative effects on non-target soil nematodes, earthworms, microarthropods etc.

Biological control with *Duddingtonia flagrans* is applicable to the larvae of: SEE LABEL FOR FULL LIST

Sheep & Goats: Barber's Pole Worm or Wire Worm (*Haemonchus* spp.), Black Scour Worm or Hair Worm (*Trichostrongylus* spp.), Brown Stomach Worm (*Teladorsagia* (*Ostertagia*) spp.), Nodule Worm (*Oesophagostomum* spp.), Thin-necked Intestinal Worm (*Nematodirus* spp.) and Hookworm (*Bunostomum* spp.).

Cattle: Barber's Pole Worm or Wire Worm (*Haemonchus* spp.), Brown Stomach Worm (*Ostertagia* spp.), Black Scour Worm or Hair Worm (*Trichostrongylus* spp.), Hookworm (*Bunostomum* spp.), Intestinal Worm (*Cooperia* spp.), Thin-necked Intestinal Worm (*Nematodirus* spp.), Nodule Worm (*Oesophagostomum* spp.).

Horses: Large strongyles (large red worms), including *Strongylus* spp., *Triodontophorus* spp. and *Oesophagodontus* spp., small strongyles (small red worms or cyathostomes), including *Cyathostomum* spp., *Cylicocyclus* spp. and *Cylicostephanus* spp., Stomach Hair Worm (*Trichostrongylus axei*), Ascarids (*Parascaris equorum*), Threadworms (*Strongyloides westeri*) and Pinworms (*Oxyuris equi*).

Other grazing animals: including Deer, Alpacas and zoo animals

DOSAGE AND ADMINISTRATION: **Abridged** see **DIRECTIONS FOR USE** on label and read carefully

1. Treat animals with a suitable chemical wormer.
 2. Move treated animals onto low worm pasture (ideally not grazed by the same animal species for a minimum 6 weeks).
 3. Commence daily use of *Livamol*[®] with *BioWorma*[®] to minimise pasture infectivity and maintain the animal's low worm status.
 4. Thoroughly mix *Livamol*[®] with *BioWorma*[®] with feed or feed supplements. *Livamol*[®] with *BioWorma*[®] will begin to work immediately within the manure.
 5. Recommended for strategic use during periods when weather conditions are conducive to larval development and transmission on pasture at temperatures above 5° Celsius (40° Fahrenheit) throughout the year.
- Use in conjunction with a recommended worm management strategy program for your area.

DAILY FEEDING RATES

LIVAMOL[®] with *BioWorma*[®] may be mixed with feed, offered by free access alongside other feed or may be used in formulated rations preferably under the direction of a nutritionist or veterinarian.

(A standard coffee mug (300 mL) holds 200 g *LIVAMOL*[®] with *BioWorma*[®])

Bodyweight * (kg)	25	50	100	200	300	400	500**
Dosage (g/head/day)	25	50	100	200	300	400	500

*Dose according to heaviest animal in the group. ** Use additional 100 g for each 100 kg thereafter.

WITHHOLDING PERIOD: MEAT: Zero (0) days, MILK: NIL Zero (0) days.

POISONS SCHEDULE: NOT APPLICABLE (Australia)

REGULATORY STATUS: Australia: APVMA No. 82646

New Zealand: ACVM No. A011335

PACK SIZES: 7.5 kg, 10 kg, 15 kg, 20 kg

Australia-Wide Toll Free: 1800 801 201	New Zealand Toll Free: 0800 424 999 (0800 IAH 999)
Email: info@iahp.com.au	Website: www.iahp.com.au

© *LIVAMOL*, *BioWorma* and the Australia/World symbol are registered trademarks of International Animal Health Products P/L.

© Copyright 2018. All Rights Reserved. Patent Pending

04/2018